

January 1994

Chiral symmetry breaking in the $d = 3$ Nambu-Jona-Lasinio model in curved spacetime

E. ELIZALDE ¹

Department E.C.M. and I.F.A.E., Faculty of Physics, University of Barcelona,
 Diagonal 647, 08028 Barcelona,
 and Center for Advanced Studies, C.S.I.C., Cam de Santa Brbara,
 17300 Blanes, Catalonia, Spain

S.D. ODINTSOV ²

Department E.C.M., Faculty of Physics, University of Barcelona,
 Diagonal 647, 08028 Barcelona, Catalonia, Spain
 and YU.I. SHIL'NOV

Department of Theoretical Physics, Kharkov State University,
 Svobody sq. 4, Kharkov 310077, Ukraine

Abstract

The phase structure of the $d = 3$ Nambu-Jona-Lasinio model in curved spacetime is considered to leading order in the $1/N$ -expansion and in the linear curvature approximation. The possibility of a curvature-induced first-order phase transition is investigated numerically. The dynamically generated fermionic mass is calculated for some values of the curvature.

¹E-mail: eli@ebubecm1.bitnet, eli@zeta.ecm.ub.es

²E-mail: odintsov@ebubecm1.bitnet; on leave from Tomsk Pedagogical Institute, 634041 Tomsk, Russian Federation.

1. Four-fermion models provide a very useful laboratory for the analytical study of composite bound states and dynamical symmetry breaking. A particularly interesting theory belonging to this class is the $d = 3$ four-fermion model, which is the natural $d = 3$ analog of the well-known Nambu-Jona-Lasinio model [1] (for a recent discussion, in connection with dynamical symmetry breaking in electroweak interactions, see [2]). This model is known to be renormalizable in the $1/N$ -expansion [3], and has a quite interesting phase structure in $d = 3$ flat space with non-trivial topology [4]. From a different point of view, the $d = 3$ four-fermion model can be considered as the high-temperature phase of the corresponding $d = 4$ model, what gives an additional motivation for its study. Let us also recall that four-fermion models are interesting, in general, as effective theories of QCD (for a comprehensible introduction to QCD, see [5]).

Recently, a careful investigation of the $d = 4$ four-fermion theory (i.e., the Nambu-Jona-Lasinio model [1]) in curved spacetime has been undertaken [6]-[8]. In particular, it has been shown in detail the existence of a first-order phase transition —induced by the curvature— from the chiral symmetric phase to the chiral non-symmetric one [7, 8]. The dynamical generation of the fermion mass due to curvature effects has also been discussed —a process that may have interesting applications in the very early universe (see, for example, [9]).

In the present note we will investigate the $d = 3$ four-fermion model in curved spacetime. The effective potential for the composite field $\bar{\psi}\psi$ will be calculated in the $1/N$ -expansion and its phase structure will be discussed. In particular, the possibility of chiral symmetry breaking and the influence of curvature in this phenomenon will be studied.

2. Our starting point is the Lagrangian of the $d = 3$ four-fermion model in curved spacetime:

$$L = \bar{\psi}^i \gamma^\mu(x) \nabla_\mu \psi^i - \frac{N\sigma^2}{2\lambda} - \sigma \bar{\psi}^i \psi_i, \quad (1)$$

where $i = 1, \dots, N$, and N is the number of fermion species. The Lagrangian (1) is equivalent to the four-fermion model, what is easily seen by solving for the equation which involves the composite field σ . Working in the linear curvature approximation and using the technique described in detail in [7] —for the $d = 4$ Nambu-Jona-Lasinio model— we obtain, after some calculations, the effective potential for the composite field σ to leading order in the $1/N$ -expansion (we shall spare the reader the details of the rather tedious computation)

$$V(\sigma) = \frac{\sigma^2}{2\lambda} - \frac{1}{3\pi^2} \left[\Lambda^3 \ln \left(1 + \frac{\sigma^2}{\Lambda^2} \right) + 2\sigma^2 \left(\Lambda - \sigma \arctan \frac{\Lambda}{\sigma} \right) \right] - \frac{R}{36\pi^2} \left(\frac{2\sigma^2\Lambda}{\Lambda^2 + \sigma^2} - 3\sigma \arctan \frac{\Lambda}{\sigma} \right). \quad (2)$$

Here Λ is a cut-off, and the curvature is considered to be arbitrary, but small, so that it is

meaningful to keep only linear curvature corrections to the flat potential, dropping $\mathcal{O}(R^2)$ -terms and those of higher order.

We now start with the analysis of Eq. (2). Consider first the case $R = 0$. Then, if $\lambda < \lambda_c = \pi^2/(2\Lambda)$, it turns out that $\sigma = 0$ is the minimum of the potential and the theory is in its chiral symmetric phase. If, on the other hand, $\lambda > \lambda_c = \pi^2/(2\Lambda)$, then $\sigma = 0$ is a maximum, and the minimum of the effective potential is now given by the solution of the equation

$$\frac{\sigma}{\Lambda} \arctan \frac{\Lambda}{\sigma} = 1 - \frac{\lambda_c}{\lambda}. \quad (3)$$

In this region the chiral symmetry of the classical Lagrangian is broken. One may ask for finiteness of σ as $\Lambda \rightarrow \infty$. This can be obtained by means of the corresponding standard renormalization of λ :

$$\lambda = \frac{\lambda_c \frac{\Lambda}{\sigma}}{\frac{\Lambda}{\sigma} - \arctan \frac{\Lambda}{\sigma}}. \quad (4)$$

The corresponding “ β -function” is defined as (see, for instance, Ref. [10])

$$\beta = \lim_{\Lambda \rightarrow \infty} \frac{\partial \lambda}{\partial \log \Lambda} = \frac{\lambda}{\lambda_c} (\lambda_c - \lambda). \quad (5)$$

Such a β -function indicates that λ_c is an ultraviolet stable fixed point.

3. Let us now proceed with the numerical computation of the effects of curvature on chiral symmetry breaking. The conditions for a first-order phase transition are given by

$$V(\sigma)|_{\sigma=\sigma_0} = 0, \quad \left. \frac{\partial V(\sigma)}{\partial \sigma} \right|_{\sigma=\sigma_0} = 0, \quad (6)$$

which yield a minimum of $V(\sigma)$. They can be written as: $y(t) = 0$, $y'(t) = 0$, namely

$$\begin{aligned} \frac{t^2}{x} - \frac{1}{3} \left[\ln(1+t^2) + 2t^2 \left(1 - t \arctan \frac{1}{t} \right) \right] - \frac{\rho}{36} \left(\frac{2t^2}{1+t^2} - 3t \arctan \frac{1}{t} \right) &= 0, \\ \frac{2t}{x} - 2t \left(1 - t \arctan \frac{1}{t} \right) - \frac{\rho}{36} \left[\frac{4t}{(1+t^2)^2} + \frac{3t}{1+t^2} - 3 \arctan \frac{1}{t} \right] &= 0 \end{aligned} \quad (7)$$

where

$$t \equiv \frac{\sigma}{\Lambda}, \quad y \equiv \frac{V}{\Lambda^3}, \quad x \equiv \frac{\lambda}{\lambda_c}, \quad \rho \equiv \frac{R}{\Lambda^2}. \quad (8)$$

It can be checked that the solution, t_0 , of these two equations readily satisfies $y''(t_0) > 0$ (for $x > 1$), with

$$y''(t) = \frac{2}{\pi^2} \left\{ \frac{1}{x} - 2 + \frac{1}{1+t^2} + 2t \arctan \frac{1}{t} - \frac{\rho}{36} \left[\frac{8}{(1+t^2)^3} - \frac{3}{(1+t^2)^2} \right] \right\}, \quad (9)$$

and it is thus, in fact, a minimum of $y(t)$. Numerical results, corresponding to several values of λ/λ_c , together with the respective values of the critical curvature R_c/Λ^2 in each case, are given in Table 1.

λ/λ_c	R_c/Λ^2	σ_0/Λ
1.03	0.00158	0.014
1.10	0.016	0.046
1.27	0.10	0.12
2.00	0.80	0.39

Table 1: Numerical values of the dynamical fermionic mass σ_0/Λ at the point of the curvature-induced phase transition —together with the respective values of the critical curvature R_c/Λ^2 —corresponding to a sample of values of $x \equiv \lambda/\lambda_c > 1$.

Table 1 shows how the dynamically generated fermionic mass changes in terms of the value of the corresponding critical curvature, for several particular values of $\lambda > \lambda_c$. Notice that in all the cases considered, the critical curvature is always smaller than 1, and also clearly $R_c/\Lambda^2 < \sigma_0/\Lambda$ (this improves for λ approaching λ_c), while R_c is of the order of σ_0^2 . However, observe also that while this last bound improves when λ departs from λ_c , $\lambda > \lambda_c$, then the value of R_c/Λ^2 approaches 1. One reaches a compromise for

$$1.1 < \frac{\lambda}{\lambda_c} < 1.5,$$

where all the requirements of the theory are met. The results are given in Figs. 1-3, where the values of V/Λ^3 as a function of σ/Λ are depicted, for $x \equiv \lambda/\lambda_c = 1.272$ and R_c/Λ^2 equal to 0.05 (Fig. 1), 0.10 (Fig. 2), and 0.15 (Fig. 3), respectively. Finally, Fig. 4 is a surface plot of the effective potential V/Λ^3 as a function of both σ/Λ and R/Λ^2 , in the intervals $[0, 0.2]$ and $[0.05, 0.15]$, respectively, for fixed $x \equiv \lambda/\lambda_c = 1.272$. Notice that Figs. 1-3 correspond to the extreme and middle xz -sections of this surface plot.

4. To summarize, we have here investigated the influence of curvature to the chiral symmetry breaking pattern of the $d = 3$ Nambu-Jona-Lasinio model. As happens in the case of the four-dimensional model [7] (and within the approximation at which we are working), we have shown that a first-order phase transition induced by curvature does also take place for $d = 3$. Of course, this is a result obtained by using the linear curvature approximation. It would be of interest to study the same model in the strong curvature phase, for instance, considering it immersed in a general De Sitter space. Then, the phase structure at arbitrarily high curvature could be investigated. There exists thus the possibility that chiral symmetry could be broken even in the region $\lambda < \lambda_c$ —where the corresponding theory in flat space is in its chiral symmetric phase.

Acknowledgments

SDO would like to thank T. Muta and T. Inagaki for helpful discussions, and the members of the Dept. ECM, Barcelona University, for their kind hospitality. This work has been supported by DGICYT (Spain) and by CIRIT (Generalitat de Catalunya).

References

- [1] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961).
- [2] Y. Nambu, in *Proceedings of the International Workshop on New Trends in Strong Coupling Gauge Theories*, Eds. M. Bando, T. Muta and K. Yamawaki (Nagoya, 1988); V.A. Miransky, M. Tanabashi and K. Yamawaki, Phys. Lett. **B221**, 177 (1989); W.J. Marciano, Phys. Rev. Lett. **62**, 2793 (1989); W.A. Bardeen, C.T. Hill and M. Lindner, Phys. Rev. **D41**, 1647 (1990).
- [3] B. Rosenstein, B.J. Warr and S.H. Park, Phys. Rep. **205**, 59 (1991).
- [4] D.Y. Song, Phys. Rev. **D48**, 3925 (1993).
- [5] T. Muta, *Foundations of Quantum Chromodynamics* (World Sci., Singapore, 1987).
- [6] C. Hill and D.S. Salopek, Ann. Phys. (NY) **213**, 21 (1992); T. Muta and S.D. Odintsov, Mod. Phys. Lett. **A6**, 3641 (1991).
- [7] T. Inagaki, T. Muta and S.D. Odintsov, Mod. Phys. Lett. **A8**, 2117 (1993).
- [8] E. Elizalde, S. Leseduardte and S.D. Odintsov, preprint hep-th 9312164.
- [9] I.L. Buchbinder, S.D. Odintsov and I.L. Shapiro, *Effective Action in Quantum Gravity* (IOP Publishing, Bristol, 1992).
- [10] K.-I. Kondo, M. Tanabashi and K. Yamawaki, Nagoya preprint DPNU-92-93, Progr. Theor. Phys., to appear.

Figure captions

Figure 1. Plot of the effective potential V/Λ^3 as a function of σ/Λ in the interval $[0, 0.2]$, for fixed values of $x \equiv \lambda/\lambda_c = 1.272$ and $R/\Lambda^2 = 0.05$.

Figure 2. The effective potential V/Λ^3 as a function of σ/Λ in the interval $[0, 0.2]$, for the same $x \equiv \lambda/\lambda_c = 1.272$, but now $R/\Lambda^2 \equiv R_c/\Lambda^2 = 0.10$. The numerical value of the dynamically generated fermionic mass $\sigma_0/\Lambda = 0.12$ can be explicitly taken from the plot in this case.

Figure 3. Plot of V/Λ^3 as a function of σ/Λ in the interval $[0, 0.2]$, again for $x \equiv \lambda/\lambda_c = 1.272$, but now $R/\Lambda^2 = 0.15$.

Figure 4. Surface plot of V/Λ^3 as a function of both σ/Λ and R/Λ^2 , in the intervals $[0, 0.2]$ and $[0.05, 0.15]$, respectively, for fixed $x \equiv \lambda/\lambda_c = 1.272$. Figs. 1-3 correspond to the extreme and middle xz -sections of this plot.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9401056v1>

This figure "fig2-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9401056v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9401056v1>

This figure "fig2-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9401056v1>